

## APCalcBC-HomeworkQuiz-#1


1. If  $f$  and  $g$  are twice differentiable functions such that  $g(x) = e^{f(x)}$  and  $g''(x) = h(x)e^{f(x)}$ , then  $h(x) =$

- (A)  $f'(x) + f''(x)$   
 (B)  $f'(x) + (f''(x))^2$   
 (C)  $(f'(x) + f''(x))^2$   
 (D)  $(f'(x))^2 + f''(x)$   
 (E)  $2f'(x) + f''(x)$


$$\begin{aligned} g(x) &= e^{f(x)} \\ g'(x) &= e^{f(x)} \cdot f'(x) \\ g''(x) &= e^{f(x)} f''(x) + f'(x) e^{f(x)} f'(x) \\ &= [f''(x) + (f'(x))^2] e^{f(x)} \end{aligned}$$

2. Let  $f$  be a function that is continuous on the closed interval  $[2, 4]$  with  $f(2) = 10$  and  $f(4) = 20$ . Which of the following is guaranteed by the Intermediate Value Theorem?

- (A)  $f(x) = 13$  has at least one solution in the open interval  $(2, 4)$ .  
 (B)  $f(3) = 15$   
 (C)  $f$  attains a maximum on the open interval  $(2, 4)$ .  
 (D)  $f'(x) = 5$  has at least one solution in the open interval  $(2, 4)$ .  
 (E)  $f'(x) > 0$  for all  $x$  in the open interval  $(2, 4)$ .



3.



Graph of  $f$

The graph of the function  $f$  is shown in the figure above. The value of  $\lim_{x \rightarrow 1^+} f(x)$  is

## APCalcBC-HomeworkQuiz-#1

- (A)  $-2$   
 (B)  $-1$   
 (C)  $2$   
 (D) nonexistent

4. A person 2 meters tall walks directly away from a streetlight that is 8 meters above the ground. If the person is walking at a constant rate and the person's shadow is lengthening at the rate of  $\frac{4}{9}$  meter per second, at what rate, in meters per second, is the person walking?

- (A)  $\frac{4}{27}$   
 (B)  $\frac{4}{9}$   
 (C)  $\frac{3}{4}$   
 (D)  $\frac{4}{3}$   
 (E)  $\frac{16}{9}$



$$\frac{ds}{dt} = \frac{4}{9}$$


$$\frac{x+s}{s} = \frac{8}{2} = 4$$

$$x+s = 4s$$

$$x = 3s$$

$$\frac{dx}{dt} = 3 \frac{ds}{dt} = 3 \left( \frac{4}{9} \right) = \frac{12}{9} = \frac{4}{3}$$

5.



Left

Right

midpt


(lowest)

trapezoidal

The graph of the function  $f$  is shown above for  $0 \leq x \leq 3$ . Of the following, which has the least value?

- (A)  $\int_1^3 f(x) dx$   
 (B) Left Riemann sum approximation of  $\int_1^3 f(x) dx$  with 4 subintervals of equal length  
 (C) Right Riemann sum approximation of  $\int_1^3 f(x) dx$  with 4 subintervals of equal length  
 (D) Midpoint Riemann sum approximation of  $\int_1^3 f(x) dx$  with 4 subintervals of equal length  
 (E) Trapezoidal sum approximation of  $\int_1^3 f(x) dx$  with 4 subintervals of equal length

## APCalcBC-HomeworkQuiz-#1



$$L = 200$$

$$\frac{dP}{dt} = kP \left(1 - \frac{P}{L}\right) = kP - \frac{k}{L}P^2$$

↑  
must be negative

Which of the following differential equations for a population  $P$  could model the logistic growth shown in the figure above?

- (A)  $\frac{dP}{dt} = 0.2P - 0.001P^2$   
 (B)  $\frac{dP}{dt} = 0.1P - 0.001P^2$   
 (C)  $\frac{dP}{dt} = 0.2P^2 - 0.001P$   
 (D)  $\frac{dP}{dt} = 0.1P^2 - 0.001P$   
 (E)  $\frac{dP}{dt} = 0.1P^2 + 0.001P$  (not this)

$$\frac{dP}{dt} = kP - \frac{k}{L}P^2$$

$$\frac{k}{L} = 0.001, L = 200$$

$$\text{so } \frac{k}{200} = 0.001, k = 0.2$$

$$\frac{dP}{dt} = 0.2P - 0.001P^2$$

7.  $\lim_{x \rightarrow 0} \frac{e^x - \cos x - 2x}{x^2 - 2x}$  is

(A)  $-\frac{1}{2}$

(B) 0

(C)  $\frac{1}{2}$

(D) 1

(E) nonexistent

$$\frac{1-1-0}{0-0} = \frac{0}{0}$$

$$\text{L'Hop} = \lim_{x \rightarrow 0} \frac{e^x + \sin x - 2}{2x - 2} = \frac{1+0-2}{0-2} = \frac{-1}{-2} = \frac{1}{2}$$

8. A curve is defined by the parametric equations  $x(t) = t^2 + 3$  and  $y(t) = \sin(t^2)$ . Which of the following is an expression for  $\frac{d^2y}{dx^2}$  in terms of  $t$ ?

(A)  $-\sin(t^2)$

(B)  $-2t \sin(t^2)$

(C)  $\cos(t^2) - 2t^2 \sin(t^2)$

(D)  $2 \cos(t^2) - 4t^2 \sin(t^2)$

$$\frac{dy}{dx} = \frac{\cos(t^2) 2t}{2t} = \cos(t^2)$$

$$\frac{d^2y}{dx^2} = \frac{\frac{d}{dt} \left( \frac{dy}{dx} \right)}{\left( \frac{dx}{dt} \right)} = \frac{-\sin(t^2) 2t}{2t} = -\sin(t^2)$$

9. Let  $f(x) = \int_{-2}^{x^2-3x} e^{t^2} dt$ . At what value of  $x$  is  $f(x)$  a minimum?

## APCalcBC-HomeworkQuiz-#1

(A) For no value of  $x$ (B)  $\frac{1}{2}$ (C)  $\frac{3}{2}$ 

(D) 2

(E) 3

$$f(x) = \int_{-2}^{x^2-3x} e^{t^2} dt \quad f(x) \text{ min when } f'(x) = 0$$

$$f'(x) = \frac{d}{dx} \left[ \int_{-2}^{x^2-3x} e^{t^2} dt \right] = e^{(x^2-3x)^2} (2x-3) = 0$$

$$\underline{\hspace{1cm}} \quad x = 3/2$$

10. The third-degree Taylor polynomial about  $x = 0$  of  $\ln(1-x)$  is

(A)  $-x - \frac{x^2}{2} - \frac{x^3}{3}$

(B)  $1 - x + \frac{x^2}{2}$

(C)  $x - \frac{x^2}{2} + \frac{x^3}{3}$

(D)  $-1 + x - \frac{x^2}{2}$

(E)  $-x + \frac{x^2}{2} - \frac{x^3}{3}$

$$P_3 = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3$$

$$f(x) = \ln(1-x) \quad f(0) = \ln 1 = 0$$

$$f'(x) = \frac{1}{1-x} (-1) \quad f'(0) = \frac{-1}{1-0} = -1$$

$$f''(x) = \frac{(1-x)(0) - (-1)(-1)}{(1-x)^2} = \frac{-1}{(1-x)^2} \quad f''(0) = \frac{-1}{1} = -1$$

$$f'''(x) = \frac{(1-x)^3(0) - (-1)(2(1-x)(-1))}{(1-x)^4}$$

$$= \frac{-2(1-x)}{(1-x)^4} = \frac{-2}{(1-x)^3} \quad f'''(0) = \frac{-2}{1} = -2$$

$$P_3 = 0 + (-1)x + \frac{(-1)}{2}x^2 + \frac{-2}{6}x^3$$

$$= -x - \frac{1}{2}x^2 - \frac{1}{3}x^3$$